

ON THE TOPOLOGY OF MANIFOLDS WITH POSITIVE ISOTROPIC CURVATURE

SIDDARTHA GADGIL AND HARISH SESHADRI

ABSTRACT. We show that a closed orientable Riemannian n -manifold, $n \geq 5$, with positive isotropic curvature and free fundamental group is homeomorphic to the connected sum of copies of $S^{n-1} \times S^1$.

1. INTRODUCTION

Let (M, g) be a closed, orientable, Riemannian manifold with positive isotropic curvature. By [9], if M is simply-connected then M is homeomorphic to a sphere of the same dimension. We shall generalise this to the case when the fundamental group of M is a free group.

Theorem 1.1. *Let M be a closed, orientable Riemannian n -manifold with positive isotropic curvature. Suppose that $\pi_1(M)$ is a free group on k generators. Then, if $n \neq 4$ or $k = 1$ (i.e. $\pi_1(M) = \mathbb{Z}$), M is homeomorphic to the connected sum of k copies of $S^{n-1} \times S^1$.*

We note that a conjecture of M. Gromov ([4] Section 3 (b)) and A. Fraser [2], based on the work of Micallef-Wang [8], states that any compact manifold with positive isotropic curvature has a finite cover satisfying our hypothesis.

Conjecture 1 (M. Gromov-A. Fraser). *$\pi_1(M)$ is virtually free, i.e., it is a finite extension of a free group.*

It is known by the work of A. Fraser [2] and A. Fraser and J. Wolfson [3] that $\pi_1(M)$ does not contain any subgroup isomorphic to the fundamental group of a closed surface of genus at least one.

Our starting point is the following fundamental result of M. Micallef and J. Moore [9].

Theorem 1.2 (M. Micallef-J. Moore). *Suppose M is a closed manifold with positive isotropic curvature. Then $\pi_i(M) = 0$ for $2 \leq i \leq \frac{n}{2}$.*

It is clear that the following purely topological result, together with the Micallef-Moore theorem, implies Theorem 1.1

Theorem 1.3. *Let M be a smooth, orientable, closed n -manifold such that $\pi_1(M)$ is a free group on k generators and $\pi_i(M) = 0$ for $2 \leq i \leq \frac{n}{2}$. If $n \neq 4$ or $k = 1$, then M is homeomorphic to the connected sum of k copies of $S^{n-1} \times S^1$.*

1991 *Mathematics Subject Classification.* Mathematics Subject Classification (1991): Primary 53C21, Secondary 53C20.

Henceforth let M be a smooth, orientable, closed n -manifold such that $\pi_1(M)$ is a free group on k generators and $\pi_i(M) = 0$ for $2 \leq i \leq \frac{n}{2}$. We assume throughout that all manifolds we consider are orientable.

Let $\widetilde{M}$ be the universal cover of M . Hence $\pi_1(\widetilde{M})$ is trivial and so is $\pi_i(\widetilde{M}) = \pi_i(M)$ for $2 \leq i \leq \frac{n}{2}$. We shall show that the homology of $\widetilde{M}$ is isomorphic as $\pi_1(M)$ -modules to that of the connected sum of k copies of $S^{n-1} \times S^1$. We then show that M is homotopy equivalent to the connected sum of k copies of $S^{n-1} \times S^1$ using Theorems of Whitehead. Finally, recent results of Kreck and Lück allow us to conclude the result.

Acknowledgements. We thank the referees for helpful comments and for suggesting a simplification of our proof.

2. THE HOMOLOGY OF $\widetilde{M}$

Let X denote the wedge $\vee_{j=1}^k S^1$ of k circles and let x denote the common point on the circles. Choose and fix an isomorphism φ from $\pi_1(M, p)$ to $\pi_1(X, x)$ for some basepoint $p \in M$. We shall use this identification throughout. Denote $\pi_1(M, p) = \pi_1(X, x)$ by π .

As X is an Eilenberg-MacLane space, there is a map $f : (M, p) \rightarrow (X, x)$ inducing φ on fundamental groups and a map $s : (X, x) \rightarrow (M, p)$ so that $f \circ s : X \rightarrow X$ is homotopic to the identity.

We deduce the homology of $\widetilde{M}$ using the Hurewicz Theorem and Poincaré duality.

Lemma 2.1. *For $1 \leq i \leq n/2$, $H_i(\widetilde{M}, \mathbb{Z}) = 0$*

Proof. As $\widetilde{M}$ is simply-connected and $\pi_i(\widetilde{M}) = \pi_i(M) = 0$ for $1 < i \leq n/2$ (by hypothesis), by Hurewicz theorem $H_i(\widetilde{M}, \mathbb{Z}) = 0$ for $1 \leq i \leq n/2$. $\square$

We deduce the homology in dimensions above $n/2$ using Poincaré duality for M with coefficients in the module $\mathbb{Z}[\pi]$, namely

$$H_{n-i}(M, \mathbb{Z}[\pi]) = H^i(M, \mathbb{Z}[\pi])$$

Recall that $H_k(M, \mathbb{Z}[\pi]) = H_k(\widetilde{M}, \mathbb{Z})$ and the group $H^i(M, \mathbb{Z}[\pi])$ is the cohomology with compact support $H_c^i(\widetilde{M}, \mathbb{Z})$. Hence Poincaré duality with coefficients in $\mathbb{Z}[\pi]$ is the same as Poincaré duality for a non-compact manifold relating homology to cohomology with compact support.

To apply Poincaré duality, we need the following lemma.

Lemma 2.2. *For $1 \leq i \leq n/2$, the map $s : (X, x) \rightarrow (M, p)$ induces isomorphisms of modules with $s_* : H^i(M; \mathbb{Z}[\pi]) \rightarrow H^i(X; \mathbb{Z}[\pi])$.*

Proof. As the map s induces an isomorphism on homotopy groups in dimensions at most $n/2$, it induces isomorphisms on the cohomology groups with twisted coefficients. Specifically, we can add cells of dimensions $k \geq n/2 + 2$ to M to obtain an Eilenberg-MacLane space $\bar{M}$ for the group π , which is thus homotopy equivalent to X . For $i \leq n/2$ and any $\mathbb{Z}[\pi]$ -module A , it follows that

$$H_i(M, A) = H_i(\bar{M}, A) = H_i(X, A)$$

where the first equality follows as the cells added to M to obtain $\bar{M}$ are of dimension at least $n/2 + 2$ and the second as the spaces are homotopy equivalent. $\square$

By applying Poincaré duality, we obtain the following result.

Lemma 2.3. *Let M be a smooth, orientable, closed n -manifold such that $\pi_1(M)$ is a free group on k generators and $\pi_i(M) = 0$ for $2 \leq i \leq \frac{n}{2}$. Then, for the universal cover $\widetilde{M}$ of M ,*

- (1) $H_i(\widetilde{M}, \mathbb{Z}) = 0$ for $1 \leq i < n - 1$
- (2) *We have an isomorphism $H_{n-1}(\widetilde{M}, \mathbb{Z}) = H_c^1(\widetilde{X}, \mathbb{Z})$, where $\widetilde{X}$ is the universal cover of X , determined by the isomorphisms $s_* : \pi_1(X, z) \rightarrow \pi_1(M, p)$ on fundamental groups.*

Proof. The statements follow from Lemmas 2.1 and 2.2 by using $H_*(\widetilde{M}, \mathbb{Z}) = H_*(M, \mathbb{Z}[\pi])$. $\square$

3. HOMOTOPY TYPE

We now show that M is homotopy equivalent to the connected sum Y of k copies of $S^{n-1} \times S^1$. Our first step is to construct a map $g : Y \rightarrow M$. We shall then show that it is a homotopy equivalence.

Note that Y has the structure of a CW-complex obtained as follows. The 1-skeleton of Y is the wedge X of k circles. Let α_i denote the i th circle with a fixed orientation.

We attach k $(n - 1)$ -cells D_j , with the j th attaching map mapping ∂D^{n-1} to the midpoint x_j of the j th circle. Finally, we attach a single n -cell Δ .

We associate to D_j an element $A_j \in \pi_{n-1}(Y, x)$. Namely, as the attaching map is constant, the j th $(n - 1)$ -cell gives an element $B_j \in \pi_{n-1}(Y, x_j)$. We consider the subarc β_j of α_j joining z_j to x in the negative direction and let A_j be obtained from B_j by the change of basepoint isomorphism using β_j .

Note that if we instead chose the arc joining z_j to x in the positive direction, then the resulting element is $-\alpha_j \cdot A_j$. By the construction of Y , it follows that the attaching map of the $(n - 1)$ -cell represents the element

$$\partial \Delta = \sum_j (A_j - \alpha_j \cdot A_j)$$

in $\pi_{n-1}(Y)$ regarded as a module over $\pi_1(Y)$. This can be seen for instance by using Poincaré duality.

We now construct the map $g : Y \rightarrow M$. Recall that we have a map $s : (X, z) \rightarrow (M, p)$ inducing the isomorphism φ^{-1} on fundamental groups. We define g on the 1-skeleton X of Y by $g|_X = s$. We henceforth identify the fundamental groups of Y and M using the isomorphism φ , i.e., $\pi_1(Y, z)$ is identified with π .

We next extend g to the n -cell of Y as follows. By Hurewicz theorem and Lemma 2.3, we have isomorphisms of π -modules $\pi_{n-1}(M, p) = H_{n-1}(\widetilde{M}, \mathbb{Z})$ and $\pi_{n-1}(Y, z) = H_{n-1}(\widetilde{M}, \mathbb{Z})$. By Lemma 2.3, each of these modules is isomorphic to $H_c^1(\widetilde{X}, \mathbb{Z})$ with the isomorphisms determined by the identifications of the fundamental groups.

Under the above isomorphisms the elements A_j correspond to elements A'_j in $\pi_{n-1}(M, p)$. Consider the element B'_j of $\pi_{n-1}(M, g(z_j))$ obtained from A'_j by the basechange map using the arc $f(\beta_j)$. We define the map g on D_j extending the constant map on its boundary to be a representative of B'_j .

As the π -modules $\pi_{n-1}(M, p)$ and $\pi_{n-1}(Y, z)$ are isomorphic, the image g of $\partial \Delta$ is homotopically trivial. Hence we can extend the map g across the cell Δ .

Lemma 3.1. *The map $g : Y \rightarrow M$ is a homotopy equivalence.*

Proof. Let $G : \tilde{Y} \rightarrow \tilde{M}$ be the induced map on the universal covers. By Lemma 2.3 applied to M and Y , we see that $H_p(\tilde{Y}) = H_p(\tilde{M}) = 0$ for $0 < p \neq n-1$ and G induces an isomorphism on H_{n-1} . Thus the map G is a homology equivalence. By a theorem of Whitehead [10], a homology equivalence between simply-connected CW-complexes is a homotopy equivalence.

It follows that G induces isomorphisms $G_* : \pi_k(\tilde{Y}) \rightarrow \pi_k(\tilde{M})$ for $k > 1$. As covering maps induce isomorphisms on higher homotopy groups, and g induces an isomorphism on π_1 , it follows that g is a weak homotopy equivalence, hence a homotopy equivalence (see [5]). $\square$

4. PROOF OF THEOREM 1.3

The rest of the proof of Theorem 1.3 is based on results of Kreck-Lück [7]. In [7], the authors define a manifold N to be a *Borel manifold* if any manifold homotopy equivalent to N is homeomorphic to N . We have shown that a manifold M satisfying the hypothesis of Theorem 1.3 is homotopy equivalent to the connected sum Y of k copies of $S^{n-1} \times S^1$. Hence it suffices to observe that Y is Borel.

By Theorem 0.13(b) of [7], the manifold $S^{n-1} \times S^1$ is Borel for $n \geq 4$. This completes the proof in the case when $\pi_1(M) = \mathbb{Z}$. Further, if $n \geq 5$, then Theorem 0.9 of [7] says that the connected sum of Borel manifolds is Borel, hence Y is Borel. This concludes the proof for $\pi_1(M)$ a free group and $n \geq 5$. $\square$

Finally, in the case when $n = 3$ by the Kneser conjecture (proved by Stallings) the manifold M is a connected sum of manifolds whose fundamental group is $\mathbb{Z}$. As M is orientable, it follows that if M is expressed as a connected sum of prime manifolds (such a decomposition exists and is unique by the Kneser-Milnor theorem), then each prime component is either $S^2 \times S^1$ or a homotopy sphere. By the Poincaré conjecture (Perelman's theorem), every homotopy 3-sphere is homeomorphic to a sphere. It follows that M is the connected sum of k copies of $S^2 \times S^1$. $\square$

REFERENCES

- [1] K. S. Brown *Cohomology of groups* Graduate Texts in Mathematics, **87** Springer-Verlag, New York, 1994.
- [2] A. M. Fraser *Fundamental groups of manifolds with positive isotropic curvature* Ann. of Math. (2) **158** (2003), no. 1, 345-354.
- [3] A. M. Fraser, J. Wolfson *The fundamental group of manifolds of positive isotropic curvature and surface groups* Duke Math. J. **133** (2006), no. 2, 325-334.
- [4] M. Gromov *Positive curvature, macroscopic dimension, spectral gaps and higher signatures* Functional analysis on the eve of the 21st century, Vol. II (New Brunswick, NJ, 1993), 1-213, Progr. Math., **132**, Birkhuser Boston, Boston, MA, 1996.
- [5] A. Hatcher, *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [6] H. Hopf *Fundamentalgruppe und zweite Bettische Gruppe* Comment. Math. Helv. **14** (1942), 257-309.
- [7] M. Kreck and W. Lück *Topological rigidity for non-aspherical manifolds* to appear in Quarterly Journal of Pure and Applied mathematics.
- [8] M. J. Micallef, M. Y. Wang *Metrics with nonnegative isotropic curvature*, Duke Math. J. **72** (1993), no. 3, 649-672.
- [9] M. J. Micallef, J. D. Moore *Minimal two-spheres and the topology of manifolds with positive curvature on totally isotropic two-planes*, Ann. of Math. (2) **127** (1988), no. 1, 199-227.
- [10] J. H. C. Whitehead *On simply connected, 4-dimensional polyhedra*. Comment. Math. Helv. **22**, (1949). 48-92.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE 560012, INDIA
E-mail address: `gadgil@math.iisc.ernet.in`

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE 560012, INDIA
E-mail address: `harish@math.iisc.ernet.in`